
COLUMBIA PITTAG Information System



OVERVIEW

THE COLUMBIA BASIN PIT TAG INFORMATION SYSTEM (PTAGIS) IS THE CENTRALIZED DATABASE FOR FISH MARKED WITH PASSIVE INTEGRATED TRANSPONDER (PIT) TAGS IN THE COLUMBIA RIVER BASIN.





CUSTOM SOFTWARE & HARDWARE

PTAGIS provides custom software to assist data contributors with collecting and submitting high-quality data. PTAGIS also designs, installs and maintains automated detection systems at many of the large hydropower dams on the

Columbia and Snake D9.1BF123456A

SEPARATION BY CODE (SBYC)

Individual PIT-tagged fish can be targeted by researchers for separation from the general population as they pass through juvenile bypasses or adult fishways. Target fish can be diverted to holding tanks for hands-on sampling, or collected for transportation by barge or truck.

PIT TAGS ARE ENCODED WITH A UNIQUE IDENTIFIER AND ALLOW A FISH IMPLANTED WITH ONE TO BE PASSIVELY DETECTED THROUGHOUT ITS LIFETIME.

THE RECORDS IN THE PTAGIS DATA WAREHOUSE DESCRIBE FOUR SEPARATE TYPES OF EVENTS SPECIFIC TO PIT-TAGGED FISH.

INTERROGATION

or through antennas.

A PIT-tagged fish may be detected at one

or more fixed automated detection sites. The time is recorded as the fish passes by



MARK/RELEASE

Each unique PIT tag is associated with a specific mark event. Species, size, condition, and other associated metadata are recorded for each fish, along with location and time of release.



RECAPTURE

Previously PIT-tagged fish may be recaptured and physically sampled subsequent to initial marking event.





can be inferred as a mortality event.

YEARS

RECOVERIES

LOCATIONS

TAGGING LOCATIONS

Mark/release, recapture, and recovery records are tied to locations in the Columbia Basin. These locations can be a stream segment, a fixed point, or a site within a hydropower facility. PIT tag data has been reported from nearly 700 different locations across the Basin.

COHO 4% 1 STEELHEAD 22%

SOCKEYE 2% 1

TAGGED FISH



In recent years, PIT tags have been used to mark and track an increasing number and diversity of other species throughout the Columbia Basin, including northern pikeminnow, shad, juvenile and adult Pacific lamprey, brook trout, and both resident and anadromous stocks of cutthroat trout.


CHINOOK 72%

MARKED FISH WITH PIT TAGS AND RELEASED THEM INTO THE COLUMBIA BASIN.

AROUND 40 DIFFERENT ORGANIZATIONS HAVE


Many of the dams on the Columbia and Snake rivers have interrogation equipment installed in the juvenile bypass systems and adult fish ladders. PTAGIS installs and maintains these systems with multiple redundant antennas and high availability computer systems to handle the large volume of fish passing through.

INTERROGATION



TRIBUTARIES

Low power interrogation systems were developed to provide automated detections in remote locations. These sites, operated by a variety of fisheries agencies, typically have fewer antennas which are installed directly in stream beds.



From 1987-2012, almost 14 million unique PIT-tagged fish have been detected at one of 250 sites, generating 150 million

DETECTIONS

detection records.

AUTOMATED INTERROGATION SYSTEMS PASSIVELY DETECT PIT-TAGGED FISH AS THEY PASS BY OR THROUGH SPECIAL ANTENNAS.



PTAGIS is a Fisheries Data Project of the Pacific States Marine Fisheries Commission